



Tuberculosis

Sarah Ringold; Cassio Lynn; Richard M. Glass

Online article and related content
current as of August 28, 2008.

JAMA. 2008;300(4):464 (doi:10.1001/jama.300.4.464)

<http://jama.ama-assn.org/cgi/content/full/300/4/464>

Supplementary material

Spanish PDF

<http://jama.ama-assn.org/cgi/content/full/300/4/464/DC1>

Correction

[Contact me if this article is corrected.](#)

Citations

[Contact me when this article is cited.](#)

Topic collections

Infectious Diseases; Bacterial Infections; Tuberculosis/ Other
Mycobacterium; JAMA Patient Page

[Contact me when new articles are published in these topic areas.](#)

Related Articles published in
the same issue

Tuberculosis Among Foreign-Born Persons in the United States
[Kevin P. Cain et al. *JAMA*. 2008;300\(4\):405.](#)

Opportunities and Challenges for HIV Care in Overlapping HIV and TB
Epidemics

[Diane V. Havlir et al. *JAMA*. 2008;300\(4\):423.](#)

Subscribe

<http://jama.com/subscribe>

Email Alerts

<http://jamaarchives.com/alerts>

Permissions

permissions@ama-assn.org

<http://pubs.ama-assn.org/misc/permissions.dtl>

Reprints/E-prints

reprints@ama-assn.org

Tuberculosis

Tuberculosis (TB) is infection with *Mycobacterium tuberculosis*, a type of bacterium. Tuberculosis infection typically occurs after repeated or prolonged exposure to the coughing of an actively infected person. Persons with HIV infection are at particularly high risk. Infection can involve any organ in the body, but the lungs are the most common site of damage. In **active infection**, there is damage to organs. In **latent infection**, the person carries the bacteria but does not have current signs of active infection. Latent TB infection is important to diagnose and treat because it can become active infection. **Multidrug-resistant TB** is resistant to (cannot be killed by) standard antibiotics. This type of TB arises from improper or incomplete treatment of TB infection or through exposure to a person infected with this type of bacterium. The July 23/30, 2008, issue of *JAMA* includes 2 articles about tuberculosis. This Patient Page is based on one previously published in the June 8, 2005, issue of *JAMA*.

SYMPTOMS

Symptoms vary depending on the patient's age and which organs are infected.

- Fever and sweating (particularly at night)
- Unexplained weight loss
- Loss of appetite
- Fatigue
- Persistent cough that may be associated with bloody sputum
- Difficulty breathing or chest pain with breathing

DIAGNOSIS

In addition to a complete medical history and physical examination, your doctor will perform a test for TB by injecting under the skin of your forearm a small amount of protein derived from TB bacteria. This TB skin test must be examined 48 to 72 hours later by a doctor or nurse who will measure any reaction at the injection site. If you have an active cough, your doctor will obtain a sample of your sputum to examine for the bacteria. Your doctor may also order a **chest x-ray** (a picture of the lungs) to look for signs of the infection or blood tests to determine whether other organs are infected.

TREATMENT

Appropriate treatment usually requires a regimen of 3 to 4 antibiotics taken daily for a minimum of 6 to 9 months. Treatment of multidrug-resistant TB requires additional medications and may last several years. Medications are given under close supervision to ensure that all doses are taken and to monitor for any adverse effects.


FOR MORE INFORMATION

- American Lung Association
www.lungusa.org
- National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention
Division of Tuberculosis Elimination
www.cdc.gov/tb

INFORM YOURSELF

To find this and previous JAMA Patient Pages, go to the Patient Page link on JAMA's Web site at www.jama.com. Many are available in English, Spanish, and French.

Source: American Lung Association



Sarah Ringold, MD, Writer

Cassio Lynn, MA, Illustrator

Richard M. Glass, MD, Editor

The JAMA Patient Page is a public service of JAMA. The information and recommendations appearing on this page are appropriate in most instances, but they are not a substitute for medical diagnosis. For specific information concerning your personal medical condition, JAMA suggests that you consult your physician. This page may be photocopied noncommercially by physicians and other health care professionals to share with patients. To purchase bulk reprints, call 312/464-0776.

